

A38

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
1 May 2003 (01.05.2003)

PCT

(10) International Publication Number  
WO 03/035050 A2(51) International Patent Classification<sup>7</sup>: A61K 31/00

(21) International Application Number: PCT/IB02/04290

(22) International Filing Date:  
20 September 2002 (20.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/323,314 20 September 2001 (20.09.2001) US(71) Applicant (for all designated States except US): AB SCI-  
ENCE [FR/FR]; 3, avenue Georges V, F-75008 Paris (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MOUSSY, Alain  
[FR/FR]; 22 bis, passage Dauphine, F-75006 Paris (FR).  
KINET, Jean-Pierre [FR/US]; 3 Hunt Road, Lexington,  
MA 02421 (US).(74) Agents: MARTIN, Jean-Jacques et al.; Cabinet Regim-  
beau, 20, rue de Chazelles, F-75847 Paris Cedex 17 (FR).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

## Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

## Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF TYROSINE KINASE INHIBITORS FOR PROMOTING HAIR GROWTH

(57) Abstract: The present invention relates to a method for promoting hair growth, preventing or minimizing hair loss comprising administering a tyrosine kinase inhibitor to a human in a need of such treatment, more particularly a non-toxic, selective and potent c-kit inhibitor. Preferably, said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

WO 03/035050 A2

WO 03/035050

PCT/IB02/04290

1

**Use of tyrosine kinase inhibitors for promoting hair growth**

5 The present invention relates to a method for promoting hair growth, preventing or minimizing hair loss comprising administering a tyrosine kinase inhibitor to a human in need of such treatment, more particularly a non-toxic, selective and potent c-kit inhibitor. Preferably, said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

10

Hairs are filamentous, keratinized structures consisting of a shaft and a root. The shaft is composed of specialized keratinocytes. The root lies within the hair follicle and comprises the germinative matrix and the keratogenous zone. The germinative matrix consists of pluripotent cells showing mitotic activity giving rise to the hair and its  
15 surrounding inner root sheath. Cells arising mitotically from this group move apically and differentiate along several different routes.

Hair growth depends on proliferation of hair follicle matrix cells. It alternates between phases of activity and rest. The anagen phase is a period of growth lasting for two to six  
20 years. During this time, the follicle is long and deep, and produces thick, well-pigmented hair. Usually, about 90% of all scalp hairs are in the anagen phase at a given time. This growth phase is followed by the catagen phase for few weeks, which corresponds to the follicle base shrinking. The resting period, or telogen phase, lasts for two to four months. In this phase, the follicle withers even further. Following the telogen phase, the next  
25 anagen phase begins, and the old hair is dislodged and falls out to make room for a new hair.

WO 03/035050

PCT/IB02/04290

2

Hair loss (alopecia areata and androgenetic alopecia) is extremely common among healthy adult humans, especially men. Indeed, hair loss affects approximately 50% of men at some point in their lives.

5 Alopecia androgenetic is an inherited condition, caused by a genetically determined sensitivity to the effects of dihydrotestosterone, which is believed to shorten the anagen phase of the hair cycle, causing miniaturisation of the follicles, and producing progressively finer hairs. Alopecia can also be induced by chemical agents or physical agents (e.g., during anti-cancer chemotherapy), and the condition also results from  
10 specific disease states and factors (emotional distress) and with increasing age. Alopecia typically is attributable to a disturbance in the hair renewal cycle leading to acceleration of the frequency of the cycles, which results in a shift in the population of follicles from the anagen phase to telogen. Ultimately, the hair follicles degenerate and a decrease in the number hairs in the affected area of the scalp or skin is observed.

15

In addition, hair loss can have a serious psychological impact. Studies have shown that hair loss can be associated with low self esteem, depression, introversion, and feelings of unattractiveness. This is reinforced by attitudes in Western society, which place great value on youthful appearance and attractiveness. Some studies have shown that based on  
20 appearance alone, men with hair loss are seen as less attractive, less assertive, less likeable, and less successful than men without hair loss.

As a result, there is real need in the development of cosmetic and clinical treatments for promoting hair growth, preventing or minimizing hair loss.

25

WO 03/035050

PCT/IB02/04290

3

As of today no treatment has been shown to be effective. Mud preparations and plant extracts have been proposed for enhancing hair growth in US 5,798,341, US 5,767,152, US 5,753,713, US 5,750,107, US 5,741,816, US 5,739,111, US 5,723,149, US 5,679,378, US 5,674,497, US 5,663,160, US 5,656,300, US 5,643,898, and US  
5 4,139,619.

2,4-diamino-6-piperidinopyrimidine 3-oxide (minoxidil) and its derivatives have been described in US 4,596,812, EP 522,964, EP 420,707, EP 459,890 and EP 519,819 for slowing hair loss. However, topical application of minoxidil and other agents is only partially effective and suffers from a number of serious side effects in many patients.

10 Other methods for treating alopecia include :

- the administration of anol, anethole and analogs with various mixture of herbs such as umbelliferae, magnoliaceae, labiatae and rutaceae (US 5,422,100),
- topical application of retinoic acid to the skin and hair (US 5,514,672),
- treatment of hair and scalp which chelating agents, gellan gum, vitamin precursors and  
15 derivatives, biotin,  $\gamma$ -linolenic acid, menthol, liposomes, various conditioners, humectants, and folic acid (US 5,523,078),
- cosmetic preparations containing solid particles of gold, silver or platinum (US 5,587,168).
- and compositions containing others compounds such as estrogens, sulfur, sulfide ions,  
20 vasodilators, inorganic selenium compounds, amino acids and protein extracts, garlic powder, brewer's yeast, grapefruit juice, acetic acid and kelp (US 5,629,002 and US 5,674,510).

Despite the above mentioned numerous approaches, effective solutions for stimulating hair growth and for preventing or minimizing hair loss continue to remain elusive.

25

WO 03/035050

PCT/IB02/04290

4

Numerous factors affect the hair growth cycle including heredity, hormonal deficiencies or imbalances, diet, stress, chemotherapy or aging. For example, it has been shown that various growth factors, steroid hormones, dermo-epithelial interaction, dermal vascularity, neuroectodermal factors, and the immune system are implicated; Stenn et al.,  
5 Dermatol. Clin., 14, 167-96 (1996) and Lindler et al., Am. J Pathol., 151, 1601-17 (1997).

More recently, alopecia areata has been considered to be a T-cell mediated autoimmune disease of the hair follicle Freyschmidt-Paul P et al, Curr Pharm Des 2001 Feb;7(3):213-30. In accordance with this mechanism, possible future therapeutic approaches include  
10 application of immunosuppressive cytokines like TGF- $\beta$  and IL-10.

Ruckert R. et al, Br J Dermatol 2000 Nov;143(5):1036-9 also observed that hair loss following skin inflammation may in part be mediated by keratinocyte (KC) apoptosis. The effects of different cytokines or other apoptosis stimulating agents such as interferon IFN- $\gamma$  or TNF- $\alpha$  on KC apoptosis have been demonstrated in vitro and in vivo.

15 Furthermore, histological investigations have shown that in C57BL mice and +/+ mice, the number of dermal mast cells in the bald areas was greater than that in controls.

Translation of these data to the human condition is rendered difficult by the complexity of the mechanisms involved in the "classical male pattern alopecia". Nevertheless, it has been found that the number of inflammatory infiltrates around the follicular infundibula  
20 of the alopecic vertices and non-alopecic occiputs of male pattern alopecia patients are significantly greater than the corresponding control value. Of interest, the number of mast cells in the widened fibrous tracts in the vertices of male pattern alopecia patients was found significantly greater than those in the adventitial fibrotic sheaths of control subjects and the non-alopecic occiputs of male pattern alopecia patients.

25

WO 03/035050

PCT/IB02/04290  
1 0 1 / 1 5 0 2 / 0 4 2 9 0

5

Mast cells (MC) are tissue elements derived from a particular subset of hematopoietic stem cells that express CD34, c-kit and CD13 antigens (Kirshenbaum et al, Blood. 94: 2333-2342, 1999 and Ishizaka et al, Curr Opin Immunol. 5: 937-43, 1993). Immature MC progenitors circulate in the bloodstream and differentiate in tissues. These differentiation and proliferation processes are under the influence of cytokines, one of utmost importance being Stem Cell Factor (SCF), also termed Kit ligand (KL), Steel factor (SL) or Mast Cell Growth Factor (MCGF). SCF receptor is encoded by the protooncogene c-kit, that belongs to type III receptor tyrosine kinase subfamily (Boissan and Arock, J Leukoc Biol. 67: 135-48, 2000). This receptor is also expressed on others hematopoietic or non hematopoietic cells. Ligation of c-kit receptor by SCF induces its dimerization followed by its transphosphorylation, leading to the recruitment and activation of various intracytoplasmic substrates. These activated substrates induce multiple intracellular signaling pathways responsible for cell proliferation and activation (Boissan and Arock, 2000). Mast cells are characterized by their heterogeneity, not only regarding tissue location and structure but also at the functional and histochemical levels (Aldenberg and Enerback., Histochem. J. 26: 587-96, 1994 ; Bradding et al. J Immunol. 155: 297-307, 1995 ; Irani et al, J Immunol. 147: 247-53, 1991 ; Miller et al, Curr Opin Immunol. 1: 637-42, 1989 and Welle et al, J Leukoc Biol. 61: 233-45, 1997).

20 In connection with the invention, it is proposed that mast cells play a crucial role in alopecia, in that they produce a large variety of mediators categorized here into three groups:

- preformed granule-associated mediators (histamine, proteoglycans, and neutral proteases),
- 25 - lipid-derived mediators (prostaglandins, thromboxanes and leucotrienes),
- and various cytokines (IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8, TNF- $\alpha$ , GM-CSF, MIP-1a, MIP-1b and IFN- $\gamma$ ).

WO 03/035050

PCT/IB02/04290

6

Then, liberation by activated mast cells of mediators (TNF-  $\alpha$ , leucotrienes, prostaglandines etc...) can induce inflammation around follicles leading to apoptosis of cells in the germinative matrix. This support that the inflammatory process mediated by mast cells is, at least in part, responsible for the development of alopecia. Local therapeutic strategies aiming at blocking the activation and the survival of mast cells, for instance through inhibition of c-kit or c-kit signaling can thus be beneficial and could help to decrease hair-loss in such condition.

More specifically, the present invention proposes to use c-kit specific kinase inhibitors to inhibit mast cell proliferation, survival and activation. A new route for promoting hair growth, preventing or minimizing hair loss is provided, which consists of destroying mast cells playing a role in the apoptosis of cells in the hair follicles. It has been found that tyrosine kinase inhibitors and more particularly c-kit inhibitors are especially suited to reach this goal.

### Description

The present invention relates to a method for promoting hair growth, preventing or minimizing hair loss comprising administering a tyrosine kinase inhibitor to a human in need of such treatment.

Tyrosine kinase inhibitors are selected for example from bis monocyclic, bicyclic or heterocyclic aryl compounds (WO 92/20642), vinylene-azaindole derivatives (WO 94/14808) and 1-cyclopropyl-4-pyridyl-quinolones (US 5,330,992), Styryl compounds (US 5,217,999), styryl-substituted pyridyl compounds (US 5,302,606), seleoindoles and selenides (WO 94/03427), tricyclic polyhydroxylic compounds (WO 92/21660) and

WO 03/035050

PCT/IB02/04290

7

benzylphosphonic acid compounds (WO 91/15495), pyrimidine derivatives (US 5,521,184 and WO 99/03854), indolinone derivatives and pyrrol-substituted indolinones (US 5,792,783, EP 934 931, US 5,834,504, US 5,883,116, US 5,883,113, US 5,886,020, WO 96/40116 and WO 00/38519), as well as bis monocyclic, bicyclic aryl and heteroaryl compounds (EP 584 222, US 5,656,643 and WO 92/20642), quinazoline derivatives (EP 602 851, EP 520 722, US 3,772,295 and US 4,343,940) and aryl and heteroaryl quinazoline (US 5,721,237, US 5,714,493, US 5,710,158 and WO 95/15758).

Preferably, said tyrosine kinase inhibitors are unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

In another embodiment, the invention is directed to a method for promoting hair growth, preventing or minimizing hair loss comprising administering a c-kit inhibitor to a human in need of such treatment.

Preferably, said c-kit inhibitor is a non-toxic, selective and potent c-kit inhibitor. Such inhibitors can be selected from the group consisting of indolinones, pyrimidine derivatives, pyrrolopyrimidine derivatives, quinazoline derivatives, quinoxaline derivatives, pyrazoles derivatives, bis monocyclic, bicyclic or heterocyclic aryl compounds, vinylene-azaindole derivatives and pyridyl-quinolones derivatives, styryl compounds, styryl-substituted pyridyl compounds, seleoindoles, selenides, tricyclic polyhydroxylic compounds and benzylphosphonic acid compounds.

Among preferred compounds, it is of interest to focus on pyrimidine derivatives such as N-phenyl-2-pyrimidine-amine derivatives (US 5,521,184 and WO 99/03854), indolinone derivatives and pyrrol-substituted indolinones (US 5,792,783, EP 934 931, US 5,834,504), US 5,883,116, US 5,883,113, US 5,886,020, WO 96/40116 and WO 00/38519), as well as bis monocyclic, bicyclic aryl and heteroaryl compounds (EP 584 222, US 5,656,643 and WO 92/20642), quinazoline derivatives (EP 602 851, EP 520




WO 03/035050

PCT/IB02/04290

8


722, US 3,772,295 and US 4,343,940), 4-amino-substituted quinazolines (US 3,470,182), 4-thienyl-2-(1H)-quinazolones, 6,7-dialkoxyquinazolines (US 3,800,039), aryl and heteroaryl quinazoline (US 5,721,237, US 5,714,493, US 5,710,158 and WO 95/15758), 4-anilinoquinazoline compounds (US 4,464,375), and 4-thienyl-2-(1H)-  
5 quinazolones (US 3,551,427).

So, preferably, the invention relates to a method for promoting hair growth, preventing or minimizing hair loss comprising administering a non toxic, potent and selective c-kit inhibitor which is a pyrimidine derivative, more particularly N-phenyl-2-pyrimidine-  
10 amine derivatives of formula I :



wherein the R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>13</sub> to R<sub>17</sub> groups have the meanings depicted in EP 564 409 B1, incorporated herein in the description.

15 Preferably, the N-phenyl-2-pyrimidine-amine derivative is selected from the compounds corresponding to formula II :



WO 03/035050

PCT/IB02/04290


9

Wherein R1, R2 and R3 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl or a cyclic or heterocyclic group, especially a pyridyl group;

R4, R5 and R6 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl, especially a methyl group;

and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function.

Preferably, R7 is the following group :



10

Among these compounds, the preferred are defined as follows:


R1 is a heterocyclic group, especially a pyridyl group,

R2 and R3 are H,

R4 is a C1-C3 alkyl, especially a methyl group,

15 R5 and R6 are H,

and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function, for example the group :




20 Therefore, in a preferred embodiment, the invention relates to a method for promoting hair growth, preventing or minimizing hair loss comprising the administration of an effective amount of the compound known in the art as CGP57148B :

4-(4-méthylpipérazine-1-ylméthyl)-N-[4-méthyl-3-(4-pyridine-3-yl)pyrimidine-2-ylamino]phényl]-benzamide corresponding to the following formula :

WO 03/035050

PCT/IB02/04290

10



The preparation of this compound is described in example 21 of EP 564 409 and the  $\beta$ -form, which is particularly useful is described in WO 99/03854.

5

Alternatively, the c-kit inhibitor can be selected from :

- indolinone derivatives, more particularly pyrrol-substituted indolinones,
- monocyclic, bicyclic aryl and heteroaryl compounds, quinazoline derivatives,
- and quinaxolines, such as 2-phényl-quinaxoline derivatives, for example 2-phenyl-6,7-dimethoxy quinaxoline.

10

In a preferred aspect, the invention contemplated the method mentioned above, wherein said c-kit inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

15

The expression "hair loss" refers herein to alopecia such as alopecia areata and androgenetic alopecia as well as hair loss caused by other factors. The method depicted above embraces promoting new hair growth, promoting hair growth before, during or after chemotherapy, promoting hair growth in hair transplant patients, preventing, stopping or minimizing hair fall out.

20

In a further embodiment, c-kit inhibitors as mentioned above are inhibitors of activated c-kit. In frame with the invention, the expression "activated c-kit" means a constitutively

WO 03/035050

PCT/IB02/04290

11

activated-mutant c-kit including at least one mutation selected from point mutations, deletions, insertions, but also modifications and alterations of the natural c-kit sequence (SEQ ID N°1). Such mutations, deletions, insertions, modifications and alterations can occur in the transphosphorylase domain, in the juxtamembrane domain as well as in any domain directly or indirectly responsible for c-kit activity. The expression “activated c-kit” also means herein SCF-activated c-kit. Preferred and optimal SCF concentrations for activating c-kit are comprised between  $5.10^{-7}$  M and  $5.10^{-6}$  M, preferably around  $2.10^{-6}$  M. In a preferred embodiment, the activated-mutant c-kit in step a) has at least one mutation proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants. In another preferred embodiment, the activated-mutant c-kit in step a) has a deletion in the juxtamembrane domain of c-kit. Such a deletion is for example between codon 573 and 579 called c-kit d(573-579). The point mutation V559G proximal to the juxtamembrane domain c-kit is also of interest.

15

In this regard, the invention contemplates a method for promoting hair growth, preventing or minimizing hair loss, comprising administering to a mammal in need of such treatment a compound that is a selective, potent and non toxic inhibitor of activated c-kit obtainable by a screening method which comprises :

20

- a) bringing into contact (i) activated c-kit and (ii) at least one compound to be tested; under conditions allowing the components (i) and (ii) to form a complex,
- b) selecting compounds that inhibit activated c-kit,
- c) testing and selecting a subset of compounds identified in step b), which are unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

25

This screening method can further comprise the step consisting of testing and selecting a subset of compounds identified in step b) that are inhibitors of mutant activated c-kit (for

WO 03/035050

PCT/IB02/04290

12

example in the transphosphorylase domain), which are also capable of inhibiting SCF-activated c-kit wild.

Alternatively, in step a) activated c-kit is SCF-activated c-kit wild.

- 5 A best mode for practicing this method consists of testing putative inhibitors at a concentration above 10  $\mu$ M in step a). Relevant concentrations are for example 10, 15, 20, 25, 30, 35 or 40  $\mu$ M.

- 10 In step c), IL-3 is preferably present in the culture media of IL-3 dependent cells at a concentration comprised between 0.5 and 10 ng/ml, preferably between 1 to 5 ng/ml.

Examples of IL-3 dependent cells include but are not limited to :

- cell lines naturally expressing and depending on c-kit for growth and survival. Among such cells, human mast cell lines can be established using the following procedures :

- 15 normal human mast cells can be infected by retroviral vectors containing sequences coding for a mutant c-kit comprising the c-kit signal peptide and a TAG sequence allowing to differentiate mutant c-kits from c-kit wild expressed in hematopoietic cells by means of antibodies.

- 20 This technique is advantageous because it does not induce cellular mortality and the genetic transfer is stable and gives satisfactory yields (around 20 %). Pure normal human mast cells can be routinely obtained by culturing precursor cells originating from blood obtained from human umbilical vein. In this regard, heparinated blood from umbilical vein is centrifuged on a Ficoll gradient so as to isolate mononucleated cells from other blood components. CD34<sup>+</sup> precursor cells are then purified from the isolated cells
- 25 mentioned above using the immunomagnetic selection system MACS (Miltenyi biotech). CD34<sup>+</sup> cells are then cultured at 37°C in 5 % CO<sub>2</sub> atmosphere at a concentration of 10<sup>5</sup> cells per ml in the medium MCCM ( $\alpha$ -MEM supplemented with L-glutamine, penicillin,

WO 03/035050

PCT/IB02/04290

13

streptomycin,  $5 \times 10^{-5}$  M  $\beta$ -mercaptoethanol, 20 % veal foetal serum, 1 % bovine albumin serum and 100 ng/ml recombinant human SCF. The medium is changed every 5 to 7 days. The percentage of mast cells present in the culture is assessed each week, using May-Grünwal Giemsa or Toluidine blue coloration. Anti-tryptase antibodies can also be  
5 used to detect mast cells in culture. After 10 weeks of culture, a pure cellular population of mast cells (> 98 %) is obtained.

It is possible using standard procedures to prepare vectors expressing c-kit for transfecting the cell lines established as mentioned above. The cDNA of human c-kit has been described in Yarden et al., (1987) EMBO J.6 (11), 3341-3351. The coding part of  
10 c-kit (3000 bp) can be amplified by PCR and cloned, using the following oligonucleotides :

- 5'AAGAAGAGATGGTACCTCGAGGGGTGACCC3' (SEQ ID No2) sens
- 5'CTGCTTCGCGGCCGCGTTAACTCTTCTCAACCA3' (SEQ ID No3)  
antisens

15 The PCR products, digested with Not1 and Xho1, has been inserted using T4 ligase in the pFlag-CMV vector (SIGMA), which vector is digested with Not1 and Xho1 and dephosphorylated using CIP (Biolabs). The pFlag-CMV-c-kit is used to transform bacterial clone XL1-blue. The transformation of clones is verified using the following primers :

- 20
- 5'AGCTCGTTTAGTGAACCGTC3' (SEQ ID No4) sens,
  - 5'GTCAGACAAAATGATGCAAC3' (SEQ ID No5) antisens.

Directed mutagenesis is performed using relevant cassettes is performed with routine and common procedure known in the art..

The vector Migr-1 (ABC) can be used as a basis for constructing retroviral vectors used  
25 for transfecting mature mast cells. This vector is advantageous because it contains the

WO 03/035050

PCT/IB02/04290

14

sequence coding for GFP at the 3' and of an IRES. These features allow to select cells infected by the retrovirus using direct analysis with a fluorocytometer. As mentioned above, the N-terminal sequence of c-kit c-DNA can be modified so as to introduce a Flag sequence that will be useful to discriminating heterogeneous from endogenous c-kit.

5

Other IL-3 dependent cell lines that can be used include but are not limited to:

- BaF3 mouse cells expressing wild-type or mutated form of c-kit (in the juxtamembrane and in the catalytic sites) are described in Kitayama et al, (1996), Blood 88, 995-1004 and Tsujimura et al, (1999), Blood 93, 1319-1329.
- 10 - IC-2 mouse cells expressing either c-kit<sup>WT</sup> or c-kit<sup>D814Y</sup> are presented in Piao et al, (1996), Proc. Natl. Acad. Sci. USA 93, 14665-14669.

IL-3 independent cell lines are :

- HMC-1, a factor-independent cell line derived from a patient with mast cell leukemia, 15 expresses a juxtamembrane mutant c-kit polypeptide that has constitutive kinase activity (Furitsu T et al, J Clin Invest. 1993;92:1736-1744 ; Butterfield et al, Establishment of an immature mast cell line from a patient with mast cell leukemia. Leuk Res. 1988;12:345-355 and Nagata et al, Proc Natl Acad Sci U S A. 1995;92:10560-10564).
- P815 cell line (mastocytoma naturally expressing c-kit mutation at the 814 position) 20 has been described in Tsujimura et al, (1994), Blood 83, 2619-2626.

The extent to which component (ii) inhibits activated c-kit can be measured *in vitro* or *in vivo*. In case it is measured *in vivo*, cell lines expressing an activated-mutant c-kit, which has at least one mutation proximal to Y823, more particularly between amino acids 800 25 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants, are preferred.

WO 03/035050

PCT/IB02/04290

15

Example of cell lines expressing an activated-mutant c-kit are as mentioned.

In another preferred embodiment, the method further comprises the step consisting of testing and selecting compounds capable of inhibiting c-kit wild at concentration below  
5 1  $\mu$ M. This can be measured *in vitro* or *in vivo*.

In vivo testing may comprise measuring the ability of the tyrosine kinase inhibitors to alleviate hair loss symptoms in C57BL/Ka mice (cpdm/cpdm) that spontaneously develops alopecia with many of the characteristics of human alopecia (Gijbels MJ et al,  
10 Am J Pathol 1996 Mar;148(3):941-50 and Tanii T et al, Acta Derm Venereol 1985;65(1):64-6).

Therefore, compounds are identified and selected according to the method described above are potent, selective and non-toxic c-kit wild inhibitors.

15

Alternatively, the screening method as defined above can be practiced *in vitro*. In this regard, the inhibition of mutant-activated c-kit and/or c-kit wild can be measured using standard biochemical techniques such as immunoprecipitation and western blot. Preferably, the amount of c-kit phosphorylation is measured.

20

In a still further embodiment, the invention contemplates a method for promoting hair growth, preventing or minimizing hair loss as depicted above wherein the screening comprises :

a) performing a proliferation assay with cells expressing a mutant c-kit (for example in  
25 the transphosphorylase domain), which mutant is a permanent activated c-kit, with a plurality of test compounds to identify a subset of candidate compounds targeting activated c-kit, each having an  $IC_{50} < 10 \mu$ M, by measuring the extent of cell death,



WO 03/035050

PCT/IB02/04290

16

b) performing a proliferation assay with cells expressing c-kit wild said subset of candidate compounds identified in step (a), said cells being IL-3 dependent cells cultured in presence of IL-3, to identify a subset of candidate compounds targeting specifically c-kit,

- 5 c) performing a proliferation assay with cells expressing c-kit, with the subset of compounds identified in step b) and selecting a subset of candidate compounds targeting c-kit wild, each having an  $IC_{50} < 10 \mu M$ , preferably an  $IC_{50} < 1 \mu M$ , by measuring the extent of cell death.

- 10 Here, the extent of cell death can be measured by  $^3H$  thymidine incorporation, the trypan blue exclusion method or flow cytometry with propidium iodide. These are common techniques routinely practiced in the art.

- The method according to the invention includes preventing, delaying the onset of  
15 alopecia and/or promoting hair growth in human.

Therefore, the invention embraces the use of the compounds defined above to manufacture a medicament or a cosmetic composition for promoting hair growth, preventing or minimizing hair loss.

- 20 The pharmaceutical or cosmetic compositions utilized in this invention may be administered by any number of routes including oral, transdermal, subcutaneous, and topical.

- In addition to the active ingredients, these pharmaceutical compositions may contain  
25 suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may

WO 03/035050

PCT/IB02/04290

17

be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, Pa.).

5 Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

10 Pharmaceutical compositions suitable for use in the invention include compositions wherein c-kit inhibitors are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient, which ameliorates the symptoms or condition. Therapeutic efficacy and  
15 toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are  
20 preferred. As mentioned above, a tyrosine kinase inhibitor and more particularly a c-kit inhibitor according to the invention is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

The invention also concerns a pharmaceutical or cosmetic composition for topical  
25 administration comprising a tyrosine kinase inhibitor as defined above and optionally at least one compound selected from the group consisting of :

- 2,4-diamino-6-piperidinopyrimidine 3-oxide (minoxidil) and its derivatives,

WO 03/035050

PCT/IB02/04290

18

- anol, anethole and analogs with various mixture of plant extracts such as umbelliferae, magnoliaceae, labiatae and rutaceae,
- retinoic acid, chelating agents, gellan gum, vitamin precursors and derivatives, biotin,  $\gamma$ -linolenic acid, menthol, liposomes, various conditioners, humectants, folic acid,
- 5 particles of gold, silver or platinum (US 5,587,168), estrogens, sulfur, sulfide ions, vasodilators, inorganic selenium compounds, amino acids and protein extracts.

The compositions according to the invention may be presented in all forms normally used for topical application, in particular in the form of a gel, paste, ointment, cream,

10 lotion, liquid suspension aqueous, aqueous-alcoholic or, oily solutions, or dispersions of the lotion or serum type, or anhydrous or lipophilic gels, or emulsions of liquid or semi-solid consistency of the milk type, obtained by dispersing a fatty phase in an aqueous phase or vice versa, or of suspensions or emulsions of soft, semi-solid consistency of the cream or gel type, or alternatively of microemulsions, of microcapsules, of

15 microparticles or of vesicular dispersions to the ionic and/or nonionic type. These compositions are prepared according to standard methods.

The composition according to the invention comprises any ingredient commonly used in dermatology and cosmetic. It may comprise at least one ingredient selected from

20 hydrophilic or lipophilic gelling agents, hydrophilic or lipophilic active agents, preservatives, emollients, viscosity enhancing polymers, humectants, surfactants, preservatives, antioxidants, solvents, and fillers, antioxidants, solvents, perfumes, fillers, screening agents, bactericides, odor absorbers and coloring matter.

25 As oils which can be used in the invention, mineral oils (liquid paraffin), vegetable oils (liquid fraction of shea butter, sunflower oil), animal oils, synthetic oils, silicone oils (cyclomethicone) and fluorinated oils may be mentioned. Fatty alcohols, fatty acids

WO 03/035050

PCT/IB02/04290  
PCT / IB 0 2 / 0 4 2 9 0

19

(stearic acid) and waxes (paraffin, carnauba, beeswax) may also be used as fatty substances.

As emulsifiers which can be used in the invention, glycerol stearate, polysorbate 60 and  
5 the PEG-6/PEG-32/glycol stearate mixture are contemplated.

As hydrophilic gelling agents, carboxyvinyl polymers (carbomer), acrylic copolymers  
such as acrylate/alkylacrylate copolymers, polyacrylamides, polysaccharides such as  
hydroxypropylcellulose, clays and natural gums may be mentioned, and as lipophilic  
gelling agents, modified clays such as bentones, metal salts of fatty acids such as  
10 aluminum stearates and hydrophobic silica, or alternatively ethylcellulose and  
polyethylene may be mentioned.

As hydrophilic active agents, proteins or protein hydrolysates, amino acids, polyols,  
urea, allantoin, sugars and sugar derivatives, vitamins, starch and plant extracts, in  
15 particular those of Aloe vera may be used.

As lipophilic active agents, retinol (vitamin A) and its derivatives, tocopherol (vitamin  
E) and its derivatives, essential fatty acids, ceramides and essential oils may be used.  
These agents add extra moisturizing or skin softening features when utilized.

20

If desired, a known gelling agent may be added to the composition of the invention.  
Suitable gelling agents include a synthetic high molecular weight crosslinked polymer of  
acrylic acid, more specifically an acrylate/C.sub.10-30 alkyl acrylate copolymer available  
for example under the trade name CARBOMER 1342. Other suitable gelling agents  
25 include cellulose and cellulose derivatives such as dihydroxyethyl cellulose (tradename  
ULTRAGEL).

WO 03/035050

PCT/IB02/04290

20

In addition, a surfactant can be included in the composition so as to provide deeper penetration of the ingredients and of the tyrosine kinase inhibitor.

Among the contemplated ingredients, the invention embraces penetration enhancing  
5 agents selected for example from the group consisting of mineral oil, water, ethanol, triacetin, glycerin and propylene glycol; cohesion agents selected for example from the group consisting of polyisobutylene, polyvinyl acetate and polyvinyl alcohol, and thickening agents.

10 Chemical methods of enhancing topical absorption of drugs are well known in the art. For example, compounds with penetration enhancing properties include sodium lauryl sulfate (Dugard, P. H. and Sheuplein, R. J., "Effects of Ionic Surfactants on the Permeability of Human Epidermis: An Electrometric Study," J. Invest. Dermatol., V.60, pp. 263-69, 1973), lauryl amine oxide (Johnson et. al., US 4,411,893), azone  
15 (Rajadhyaksha, US 4,405,616 and 3,989,816) and decylmethyl sulfoxide (Sekura, D. L. and Scala, J., "The Percutaneous Absorption of Alkylmethyl Sulfides," Pharmacology of the Skin, Advances In Biology of Skin, (Appleton-Century Craft) V. 12, pp. 257-69, 1972). It has been observed that increasing the polarity of the head group in amphoteric molecules increases their penetration-enhancing properties but at the expense of  
20 increasing their skin irritating properties (Cooper, E. R. and Berner, B., "Interaction of Surfactants with Epidermal Tissues: Physiochemical Aspects," Surfactant Science Series, V. 16, Reiger, M. M. ed. (Marcel Dekker, Inc.) pp. 195-210, 1987).

Suitable solvents include alkyl esters of fatty acids, preferably C.sub.1-12, more  
25 preferably C.sub.3-10, alkyl esters of saturated or unsaturated fatty acids containing 8-22 carbon atoms. Particularly preferred solvents include isopropyl myristate, octyl palmitate, WIKENOL 161 (a mixture of esters), etc. Alcohols such as ethanol, propanol,

WO 03/035050

PCT/IB02/04290

21

isopropanol, propylene glycol, etc., as well as aqueous mixtures of these alcohols may also be used.

A second class of chemical enhancers are generally referred to as co-solvents. These materials are absorbed topically relatively easily, and, by a variety of mechanisms, achieve permeation enhancement for some drugs. Ethanol (Gale et. al., U.S. Pat. No. 4,615,699 and Campbell et. al., U.S. Pat. Nos. 4,460,372 and 4,379,454), dimethyl sulfoxide (US 3,740,420 and 3,743,727, and US 4,575,515), and glycerine derivatives (US 4,322,433) are a few examples of compounds which have shown an ability to enhance the absorption of various compounds.

Any formulation which allows delivery of the active compounds of the present invention to the skin, hair and hair follicles.

Therefore, the invention also contemplates a cosmetic composition comprising a tyrosine kinase inhibitor, more particularly a c-kit inhibitor, and at least one ingredient as depicted above suitable for a topical administration. Preferably, the composition is formulated for the delivery of the tyrosine kinase inhibitor to the skin, hair or hair follicles, such as a hair-conditioning composition.

20

WO 03/035050

PCT/IB02/04290

22

**CLAIMS**

5

1. A method for promoting hair growth, preventing or minimizing hair loss, comprising administering a tyrosine kinase inhibitor to a human in need of such treatment.

10

2. A method according to claim 1, wherein said tyrosine kinase inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

3. A method for promoting hair growth, preventing or minimizing hair loss, comprising administering a c-kit inhibitor to a human in need of such treatment.

15

4. A method according to claim 3, wherein said c-kit inhibitor is a non-toxic, selective and potent c-kit inhibitor.

20

5. A method according to claim 4, wherein said inhibitor is selected from the group consisting of indolinones, pyrimidine derivatives, pyrrolopyrimidine derivatives, quinazoline derivatives, quinoxaline derivatives, pyrazoles derivatives, bis monocyclic, bicyclic or heterocyclic aryl compounds, vinylene-azaindole derivatives and pyridyl-quinolones derivatives, styryl compounds, styryl-substituted pyridyl compounds, seleoindoles, selenides, tricyclic polyhydroxylic compounds and benzylphosphonic acid compounds.

25

6. A method according to claim 4, wherein said inhibitor is selected from the group consisting of :

- pyrimidine derivatives, more particularly N-phenyl-2-pyrimidine-amine derivatives.


WO 03/035050

PCT/IB02/04290


23

- indolinone derivatives, more particularly pyrrol-substituted indolinones,
- monocyclic, bicyclic aryl and heteroaryl compounds,
- and quinazoline derivatives.

- 5 7. A method according to claim 3, wherein said inhibitor is selected from the group consisting of N-phenyl-2-pyrimidine-amine derivatives having the formula II :



- 10 Wherein R1, R2 and R3 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl or a cyclic or heterocyclic group, especially a pyridyl group;  
 R4, R5 and R6 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl, especially a methyl group;  
 and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least  
 15 one basic site, such as an amino function, preferably the following group :



8. A method according to claim 7, wherein said inhibitor is the 4-(4-méthylpipérazine-1-ylméthyl)-N-[4-méthyl-3-(4-pyridine-3-yl)pyrimidine-2 ylamino]phényl]-benzamide.

20



WO 03/035050

PCT/IB02/04290

24

9. A method according to one of claims 3 to 8, wherein said c-kit inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

10. A method according to one of claims 3 to 9, wherein said c-kit inhibitor is an  
5 inhibitor of activated c-kit.

11. A method according to one of claims 3 to 10, wherein said activated c-kit inhibitor is capable of inhibiting SCF-activated c-kit.

10 12. A method according one of claims 3 to 10, wherein said inhibitor is capable of inhibiting constitutively activated-mutant c-kit.

13. A method for promoting hair growth, preventing or minimizing hair loss, comprising administering to a human in need of such treatment a compound that is a selective,  
15 potent and non toxic inhibitor of activated c-kit obtainable by a screening method which comprises :

- a) bringing into contact (i) activated c-kit and (ii) at least one compound to be tested; under conditions allowing the components (i) and (ii) to form a complex,
- b) selecting compounds that inhibit activated c-kit,
- 20 c) testing and selecting a subset of compounds identified in step b), which are unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

14. A method according to claim 13, wherein the screening method further comprises the step consisting of testing and selecting a subset of compounds identified in step b) that  
25 are inhibitors of mutant activated c-kit, which are also capable of inhibiting SCF-activated c-kit wild.

WO 03/035050

PCT/IB02/04290

25

15. A method according to claim 13, wherein activated c-kit is SCF-activated c-kit wild in step a).

16. A method according to one of claims 13 to 15, wherein putative inhibitors are tested  
5 at a concentration above 10  $\mu$ M in step a).

17. A method according to one of claims 13 to 16, wherein IL-3 is preferably present in the culture media of IL-3 dependent cells at a concentration comprised between 0.5 and 10 ng/ml, preferably between 1 to 5 ng/ml.

10

18. A method according to claim 17, wherein IL-3 dependent cells are selected from the group consisting of mast cells, transfected mast cells, BaF3, and IC-2.

19. A method according to one of claims 13 to 18, wherein the extent to which  
15 component (ii) inhibits activated c-kit is measured *in vitro* or *in vivo*.

20. A method according to one of claims 13 to 18, further comprising the step consisting of testing and selecting compounds capable of inhibiting c-kit wild at concentration below 1  $\mu$ M.

20

21. A method according to claim 20, wherein the testing is performed *in vitro* or *in vivo*.

22. A method according to one of claims 13 to 21, wherein the inhibition of mutant-activated c-kit and/or c-kit wild is measured using standard biochemical techniques such  
25 as immunoprecipitation and western blot.

23. A method according to one of claims 13 to 21, wherein the amount of c-kit phosphorylation is measured.

WO 03/035050

PCT/IB02/04290

26

24. A method according to one of claims 13 to 23, wherein identified and selected compounds are potent, selective and non-toxic c-kit wild inhibitors.

- 5 25. A method for promoting hair growth, preventing or minimizing hair loss, comprising administering to a human in need of such treatment a c-kit inhibitor obtainable by a screening method comprising :
- a) performing a proliferation assay with cells expressing a mutant c-kit (for example in the transphosphorylase domain), which mutant is a permanent activated c-kit, with a
- 10 plurality of test compounds to identify a subset of candidate compounds targeting activated c-kit, each having an  $IC_{50} < 10 \mu M$ , by measuring the extent of cell death,
- b) performing a proliferation assay with cells expressing c-kit wild said subset of candidate compounds identified in step (a), said cells being IL-3 dependent cells cultured in presence of IL-3, to identify a subset of candidate compounds targeting specifically c-
- 15 kit,
- c) performing a proliferation assay with cells expressing c-kit, with the subset of compounds identified in step b) and selecting a subset of candidate compounds targeting c-kit wild, each having an  $IC_{50} < 10 \mu M$ , preferably an  $IC_{50} < 1 \mu M$ , by measuring the extent of cell death.
- 20
26. A method according to claim 25, wherein the extent of cell death is measured by  $^3H$  thymidine incorporation, the trypan blue exclusion method or flow cytometry with propidium iodide.
- 25 27. A method according to one of claims 1 to 26 for treating alopecia such as alopecia areata and androgenetic alopecia as well as hair loss caused by other factors.

WO 03/035050

PCT/IB02/04290

27

28. A method according to one of claims 1 to 26 for promoting new hair growth, promoting hair growth before, during or after chemotherapy, promoting hair growth in hair transplant patients, preventing, stopping or minimizing hair fall out.
- 5 29. Use of a c-kit inhibitor to manufacture a medicament or a cosmetic composition for promoting hair growth, preventing or minimizing hair loss.
30. A composition suitable for oral, transdermal, subcutaneous, and topical administration comprising a tyrosine kinase inhibitor, more particularly a c-kit inhibitor
- 10 for promoting hair growth, preventing or minimizing hair loss.
31. A pharmaceutical or cosmetic composition according to claim 30, which is suitable for topical application.
- 15 32. A composition according to claim 31, which is in the form of a gel, paste, ointment, cream, lotion, liquid suspension aqueous, aqueous-alcoholic or, oily solutions, or dispersions of the lotion or serum type, or anhydrous or lipophilic gels, or emulsions of liquid or semi-solid consistency of the milk type, obtained by dispersing a fatty phase in an aqueous phase or vice versa, or of suspensions or emulsions of soft, semi-solid
- 20 consistency of the cream or gel type, or alternatively of microemulsions, of microcapsules, of microparticles or of vesicular dispersions to the ionic and/or nonionic type.
33. A composition according to claim 31, which comprises at least one ingredient
- 25 selected from hydrophilic or lipophilic gelling agents, hydrophilic or lipophilic active agents, emollients, viscosity enhancing polymers, humectants, surfactants, preservatives, antioxidants, solvents, and fillers.

WO 03/035050

PCT/IB02/04290

28

34. A composition according to one of claims 31 to 33, which is formulated for the delivery of the tyrosine kinase inhibitor to the skin, hair or hair follicles.

35. A composition according to claim 34, which is a hair-conditioning composition.

5

WO 03/035050

PCT/IB02/04290

## SEQUENCE LISTING

&lt;110&gt; AB Science

&lt;120&gt; Use of tyrosine kinase inhibitors for promoting hair growth

&lt;130&gt; D19830 NT

&lt;150&gt; US 60/323,314

&lt;151&gt; 2001-09-20

&lt;160&gt; 5

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 976

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;223&gt; Human c-kit

&lt;400&gt; 1

```

Met Arg Gly Ala Arg Gly Ala Trp Asp Phe Leu Cys Val Leu Leu Leu
 1           5           10           15

Leu Leu Arg Val Gln Thr Gly Ser Ser Gln Pro Ser Val Ser Pro Gly
          20           25           30

Glu Pro Ser Pro Pro Ser Ile His Pro Gly Lys Ser Asp Leu Ile Val
      35           40           45

Arg Val Gly Asp Glu Ile Arg Leu Leu Cys Thr Asp Pro Gly Phe Val
      50           55           60

Lys Trp Thr Phe Glu Ile Leu Asp Glu Thr Asn Glu Asn Lys Gln Asn
 65           70           75           80

Glu Trp Ile Thr Glu Lys Ala Glu Ala Thr Asn Thr Gly Lys Tyr Thr
          85           90           95

Cys Thr Asn Lys His Gly Leu Ser Asn Ser Ile Tyr Val Phe Val Arg
      100           105           110

Asp Pro Ala Lys Leu Phe Leu Val Asp Arg Ser Leu Tyr Gly Lys Glu
      115           120           125

Asp Asn Asp Thr Leu Val Arg Cys Pro Leu Thr Asp Pro Glu Val Thr
      130           135           140

Asn Tyr Ser Leu Lys Gly Cys Gln Gly Lys Pro Leu Pro Lys Asp Leu
      145           150           155           160

Arg Phe Ile Pro Asp Pro Lys Ala Gly Ile Met Ile Lys Ser Val Lys
          165           170           175

Arg Ala Tyr His Arg Leu Cys Leu His Cys Ser Val Asp Gln Glu Gly
          180           185           190

Lys Ser Val Leu Ser Glu Lys Phe Ile Leu Lys Val Arg Pro Ala Phe
      195           200           205

Lys Ala Val Pro Val Val Ser Val Ser Lys Ala Ser Tyr Leu Leu Arg
      210           215           220

```

WO 03/035050

PCT/IB02/04290

Glu Gly Glu Glu Phe Thr Val Thr Cys Thr Ile Lys Asp Val Ser Ser  
 225 230 235 240  
 Ser Val Tyr Ser Thr Trp Lys Arg Glu Asn Ser Gln Thr Lys Leu Gln  
 245 250 255  
 Glu Lys Tyr Asn Ser Trp His His Gly Asp Phe Asn Tyr Glu Arg Gln  
 260 265 270  
 Ala Thr Leu Thr Ile Ser Ser Ala Arg Val Asn Asp Ser Gly Val Phe  
 275 280 285  
 Met Cys Tyr Ala Asn Asn Thr Phe Gly Ser Ala Asn Val Thr Thr Thr  
 290 295 300  
 Leu Glu Val Val Asp Lys Gly Phe Ile Asn Ile Phe Pro Met Ile Asn  
 305 310 315 320  
 Thr Thr Val Phe Val Asn Asp Gly Glu Asn Val Asp Leu Ile Val Glu  
 325 330 335  
 Tyr Glu Ala Phe Pro Lys Pro Glu His Gln Gln Trp Ile Tyr Met Asn  
 340 345 350  
 Arg Thr Phe Thr Asp Lys Trp Glu Asp Tyr Pro Lys Ser Glu Asn Glu  
 355 360 365  
 Ser Asn Ile Arg Tyr Val Ser Glu Leu His Leu Thr Arg Leu Lys Gly  
 370 375 380  
 Thr Glu Gly Gly Thr Tyr Thr Phe Leu Val Ser Asn Ser Asp Val Asn  
 385 390 395 400  
 Ala Ala Ile Ala Phe Asn Val Tyr Val Asn Thr Lys Pro Glu Ile Leu  
 405 410 415  
 Thr Tyr Asp Arg Leu Val Asn Gly Met Leu Gln Cys Val Ala Ala Gly  
 420 425 430  
 Phe Pro Glu Pro Thr Ile Asp Trp Tyr Phe Cys Pro Gly Thr Glu Gln  
 435 440 445  
 Arg Cys Ser Ala Ser Val Leu Pro Val Asp Val Gln Thr Leu Asn Ser  
 450 455 460  
 Ser Gly Pro Pro Phe Gly Lys Leu Val Val Gln Ser Ser Ile Asp Ser  
 465 470 475 480  
 Ser Ala Phe Lys His Asn Gly Thr Val Glu Cys Lys Ala Tyr Asn Asp  
 485 490 495  
 Val Gly Lys Thr Ser Ala Tyr Phe Asn Phe Ala Phe Lys Gly Asn Asn  
 500 505 510  
 Lys Glu Gln Ile His Pro His Thr Leu Phe Thr Pro Leu Leu Ile Gly  
 515 520 525  
 Phe Val Ile Val Ala Gly Met Met Cys Ile Ile Val Met Ile Leu Thr  
 530 535 540  
 Tyr Lys Tyr Leu Gln Lys Pro Met Tyr Glu Val Gln Trp Lys Val Val  
 545 550 555 560  
 Glu Glu Ile Asn Gly Asn Asn Tyr Val Tyr Ile Asp Pro Thr Gln Leu  
 565 570 575

WO 03/035050

PCT/IB02/04290

Pro Tyr Asp His Lys Trp Glu Phe Pro Arg Asn Arg Leu Ser Phe Gly  
 580 585 590  
 Lys Thr Leu Gly Ala Gly Ala Phe Gly Lys Val Val Glu Ala Thr Ala  
 595 600 605  
 Tyr Gly Leu Ile Lys Ser Asp Ala Ala Met Thr Val Ala Val Lys Met  
 610 615 620  
 Leu Lys Pro Ser Ala His Leu Thr Glu Arg Glu Ala Leu Met Ser Glu  
 625 630 635 640  
 Leu Lys Val Leu Ser Tyr Leu Gly Asn His Met Asn Ile Val Asn Leu  
 645 650 655  
 Leu Gly Ala Cys Thr Ile Gly Gly Pro Thr Leu Val Ile Thr Glu Tyr  
 660 665 670  
 Cys Cys Tyr Gly Asp Leu Leu Asn Phe Leu Arg Arg Lys Arg Asp Ser  
 675 680 685  
 Phe Ile Cys Ser Lys Gln Glu Asp His Ala Glu Ala Ala Leu Tyr Lys  
 690 695 700  
 Asn Leu Leu His Ser Lys Glu Ser Ser Cys Ser Asp Ser Thr Asn Glu  
 705 710 715 720  
 Tyr Met Asp Met Lys Pro Gly Val Ser Tyr Val Val Pro Thr Lys Ala  
 725 730 735  
 Asp Lys Arg Arg Ser Val Arg Ile Gly Ser Tyr Ile Glu Arg Asp Val  
 740 745 750  
 Thr Pro Ala Ile Met Glu Asp Asp Glu Leu Ala Leu Asp Leu Glu Asp  
 755 760 765  
 Leu Leu Ser Phe Ser Tyr Gln Val Ala Lys Gly Met Ala Phe Leu Ala  
 770 775 780  
 Ser Lys Asn Cys Ile His Arg Asp Leu Ala Ala Arg Asn Ile Leu Leu  
 785 790 795 800  
 Thr His Gly Arg Ile Thr Lys Ile Cys Asp Phe Gly Leu Ala Arg Asp  
 805 810 815  
 Ile Lys Asn Asp Ser Asn Tyr Val Val Lys Gly Asn Ala Arg Leu Pro  
 820 825 830  
 Val Lys Trp Met Ala Pro Glu Ser Ile Phe Asn Cys Val Tyr Thr Phe  
 835 840 845  
 Glu Ser Asp Val Trp Ser Tyr Gly Ile Phe Leu Trp Glu Leu Phe Ser  
 850 855 860  
 Leu Gly Ser Ser Pro Tyr Pro Gly Met Pro Val Asp Ser Lys Phe Tyr  
 865 870 875 880  
 Lys Met Ile Lys Glu Gly Phe Arg Met Leu Ser Pro Glu His Ala Pro  
 885 890 895  
 Ala Glu Met Tyr Asp Ile Met Lys Thr Cys Trp Asp Ala Asp Pro Leu  
 900 905 910  
 Lys Arg Pro Thr Phe Lys Gln Ile Val Gln Leu Ile Glu Lys Gln Ile  
 915 920 925



WO 03/035050

PCT/IB02/04290

Ser Glu Ser Thr Asn His Ile Tyr Ser Asn Leu Ala Asn Cys Ser Pro  
930 935 940

Asn Arg Gln Lys Pro Val Val Asp His Ser Val Arg Ile Asn Ser Val  
945 950 955 960

Gly Ser Thr Ala Ser Ser Ser Gln Pro Leu Leu Val His Asp Asp Val  
965 970 975

<210> 2  
<211> 30  
<212> DNA  
<213> Hcmo sapiens

<220>  
<223> Primer

<400> 2  
aagaagagat ggtacctcga ggggtgaccc 30

<210> 3  
<211> 33  
<212> DNA  
<213> Homo sapiens

<220>  
<223> Primer

<400> 3  
ctgcttcgcg gccgcgttaa ctcttctcaa cca 33

<210> 4  
<211> 20  
<212> DNA  
<213> Homo sapiens

<220>  
<223> Primer

<400> 4  
agctcggtta gtgaaccgtc 20

<210> 5  
<211> 20  
<212> DNA  
<213> Homo sapiens

<220>  
<223> Primer

<400> 5  
gtcagacaaa atgatgcaac 20